Darreichungsform / Dosierung / Dose / Route

1 capsule


2 capsules


3 capsules


4 capsules

1 tablet


2 tablets


3 tablets


4 tablets


_ capsule/s


_ tablet/s

_ ML

Suppository

Apply to affected Area

Buccale


Dilute 1 sache in water


half a tablet

1 drop left ear


1 drop right ear

1 drop left eye

1 drop right eye

Vaginal insertion

Inhale

Inhale with Spacer

Oral syringe

Einnahmezeitpunkt / Frequenzy



Morning



Noon

Evening

Night

1 capsule every
2 hours

1 tablet every
4 hours



2 tablets every
4 hours



Morning



Noon



Evening

Night

1 capsule every
2 hours

1 tablet every
4 hours



2 tablets every
4 hours

Beachten / Precaution:

Not with alcohol

Crush

Do not drive

Not with alcohol or drugs



Take on empty stomach



Keep medication in the fridge

Keep out of reach of babies

Keep out of reach of children

If breastfeeding do not take medication


Do not crush

Do not take when pregnant

Seek medical advice before taking if pregnant

Seek medical advice


Shake

Take until finished

Take with food

Take with water

Mögliche Nebenwirkungen / Side effects

Back pain

Blurred vision



Confusion

Constipation

Coughing

Diarrhea

Dizziness

Dizzy when getting up

Fatigue



Fever

Headache

Heartburn



Insomnia

Loss of breath

Muscular pain

Nausea

Photosensitivity

Rash



Ringing in ears

Tremors

Vomiting



Weight gain



Weight loss